

WHAT IS CLAIMED IS:

1. In a method for preparing a polypeptide in
a cellular host, where the polypeptide is heterologous to
5 the host and may be expressed in low percentage amounts
of total protein, the improvement which comprises:

joining an open reading frame DNA sequence
coding for said polypeptide with a second open reading
frame DNA sequence coding for a heterologous ubiquitin,
10 to form a fusion polypeptide;

introducing the sequence coding for said fusion
polypeptide under conditions for expression in said host,
whereby said fusion polypeptide is expressed; and

isolating said fusion polypeptide to provide
15 said second polypeptide in high yield.

2. A method according to Claim 1, wherein
said host is a eukaryotic host.

3. A method according to Claim 2, wherein
20 said eukaryotic host is yeast.

4. A method according to Claim 3, wherein
said DNA sequences are under the transcriptional
25 regulatory control of a transcriptional initiation
regulatory region comprising a promoter region for a
glycolytic enzyme.

5. A method according to Claim 4, wherein
30 said transcriptional initiation regulatory region is
inducible.

6. A method according to Claim 1, where said
host is prokaryotic.

35

0991216-031601

7. A method according to Claim 6, wherein said prokaryotic host is *E. coli*.

5 8. A method according to Claim 1, wherein said DNA sequence coding for said polypeptide is 3' to said DNA sequence coding for ubiquitin in the direction of transcription.

10 9. A method according to Claim 1, wherein said DNA sequence coding for said polypeptide is 3' to said DNA sequence coding for ubiquitin in the direction of transcription.

15 10. In a method for preparing a mammalian polypeptide in a yeast host, where the polypeptide may be expressed in low percentage amounts of total protein, the improvement which comprises:

20 joining an open reading frame DNA sequence coding for said polypeptide with a second open reading frame DNA sequence coding for heterologous ubiquitin, to form a fusion polypeptide;

25 introducing the sequence coding for said fusion polypeptide under conditions for expression in said yeast, whereby said fusion polypeptide is expressed; and isolating said fusion polypeptide in high yield.

30 11. A method according to Claim 10, wherein said conditions for expression include an inducible transcriptional initiation regulatory region.

35 12. A method according to Claim 11, where said transcriptional initiation regulatory region consists essentially of a glycolytic enzyme promoter region and ADH2 control region.

09931216-081501
109880-9727E650

13. A DNA sequence coding for ubiquitin joined to a DNA sequence coding for a mammalian polypeptide.

5 14. An expression sequence including in direction of transcription, an inducible transcriptional initiation regulatory region and a DNA sequence according to Claim 13.

10 15. A polypeptide encoded for by a DNA sequence according to Claim 13.

15 16. A polypeptide according to Claim 15, wherein said mammalian polypeptide encodes for at least a portion of proinsulin.

20 17. A polypeptide according to Claim 15, wherein said mammalian polypeptide encodes for at least a portion of IGF-1 or IGF-2.

25
30
35

add B2

FOI b7D b7E b7F b7G b7H b7I b7J b7K b7L b7M b7N b7O b7P b7Q b7R b7S b7T b7U b7V b7W b7X b7Y b7Z